


# CLEARGRAZE®



ALLIGARE.COM

## A ROBUST, YET UNCOMPLICATED, PRODUCT TO IMPROVE YOUR OPERATION EFFICIENCY

Increase pasture stocking rates and vegetation area aesthetics with Alligare's Cleargraze. This herbicide is ideal for simple and selective broadleaf weed and brush control.

### PERFORMANCE HIGHLIGHTS

- Superior control of susceptible ALS resistant weeds, broad spectrum brush, and individual weed and woody plant species
- Can be used as a tank mix additive, with Gunslinger products, for additional plant removal or as a single, effective solution
- Product is allowable in areas where manure or plant material may be moved off site
- Acceptable use in 2,4-D restricted areas, with no grazing restrictions

### FOR BEST RESULTS

1. Use a nonionic surfactant at 0.25% v/v or methylated seed oil at a rate of 1% of the total spray volume.
2. Apply the spray solution to uniformly cover the foliage of the undesirable vegetation.
3. Basal applications are best suited for low density brush areas where there is not an excess of 400 stems per acre.
4. Apply to young, actively growing weeds, and use higher rates when weeds are more mature.

### USE SITES

- rangeland, fence rows, and pastures
- grazed areas in and around non-crop sites
- forest planting sites
- wildlife management areas
- non-crop areas, CRP acres
- cut stumps



**Alligare**®

A member of the ADAMA Group

# CLEARGRAZE®

ACTIVE INGREDIENTS: TRICLOPYR 3 LB/GAL / FLUROXYPYR 1 LB/GAL



## BROADLEAF WEEDS CONTROLLED

Annual broomweed	Pigweed
Bedstraw	Plantain
Morning glories	Prickly lettuce
Cocklebur	Prickly pear
Dogfennel	Puncturevine
Kochia	Ragweed
Dewberry	Sericea lespedeza

## BRUSH SPECIES CONTROLLED

Aspen	Poison oak
Birch species	Hackberry
Blackberry	Privet
Chinese tallow tree	Prunus spp.
Locust (black & honey)	Sumac
Persimmon	Willow
Poison ivy	Yucca

## MIXING INSTRUCTIONS

1. Add ½ of the required spray volume to the spray tank and start agitation.
2. Add the required amount of Alligare Cleargraze to the spray tank.
3. Add additional tank-mix herbicide partners as needed.
4. Add any additional surfactants.
5. Agitate during the final filling of the spray tank and maintain sufficient agitation during application to ensure uniformity of the spray mixture.

**ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS**

## SIGNAL WORD

CAUTION

## APPLICATION DETAILS

MOA Group 4, Synthetic Auxin  
Use Rate 1-2% v/v, 1-4 pints per acre  
See label for site and species specifics

## WORKER SAFETY

long-sleeved shirt  
long pants  
shoes plus socks  
waterproof gloves



**Alligare®**

A member of the ADAMA Group