

# WHETSTONE®



ALLIGARE.COM

## *A RELIABLE PARTNER FOR SUPERIOR, YET SELECTIVE, CONTROL OF NUMEROUS WOODY PLANTS, BROADLEAF WEEDS, AND VINES*

As a flexible, pre- and post-emergent product, Alligare's Whetstone is a tank-mix friendly formula that provides residual control of invasive and noxious vegetation in a variety of settings.

### PERFORMANCE HIGHLIGHTS:

- Broad spectrum control of 185 annual, biennial, and perennial weeds and brush
- Provides extended pre- and post-emergent control of susceptible species
- Approved for treating vegetation in areas grazed by livestock, with no restrictions
- Increases weed spectrum of broadleaf species when included in bare-ground applications
- Registered under the EPA's Reduced Risk Pesticide Initiative

### FOR BEST RESULTS:

1. Treatment should occur when weeds and woody plants are actively growing.
2. Use a higher rate, within label range, when growing conditions are less than favorable, weed foliage is tall and dense, or when long-term residual control is desired.
3. To improve efficacy, an adjuvant should be added to all spray mixtures.

### USE SITES:

rangeland /permanent grass pastures  
railroad and utility rights-of-way  
non-irrigation ditch banks  
natural areas and wildlife habitats


Conservation Reserve Program (CRP) lands  
roadsides & other unimproved turf grasses  
fencerows and storage areas  
industrial and utility plant sites



**Alligare®**  
A member of the ADAMA Group

# WHETSTONE®

ACTIVE INGREDIENTS: AMINOPYRALID 2 LB/GAL



## TARGETED WEEDS

Black locust	Knapweed	Puncturevine	Thistles
Carolina horsenettle	Kudzu	Ragweed	Tree of heaven
Curly dock	Lespedeza, annual	Spanish needle	Tropical soda apple
Horseweed	Mimosa	Teasel	Wisteria
Silverleaf nightshade	Buffalo bur	Cocklebur	Iron weed

## BROADCAST MIXING INSTRUCTIONS

1. Add ½ of the required spray volume to the spray tank and begin agitation.
2. Add the required amount of Whetstone.
3. As needed, add additional tank-mix herbicide partners.
4. Add surfactants and any additional adjuvants.
5. Agitate during the final filling of the spray tank and maintain sufficient agitation during application to ensure uniformity of the spray mixture.

**ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS**

## SIGNAL WORD

CAUTION

## APPLICATION DETAILS

MOA Group 4, Synthetic Auxin  
Use rate 3 - 7 ounces/acre

## WORKER SAFETY

long-sleeved shirt  
long pants  
shoes plus socks



**Alligare®**

A member of the ADAMA Group

092324