



IMOX Herbicide

FIFRA 2(ee) Recommendation

EPA Reg. No. 81927-66

**FOR DISTRIBUTION AND USE IN ALL REGISTERED STATES
EXCEPT TEXAS AND OREGON**

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation for Imox Herbicide is valid until December 31, 2028, or until withdrawn, canceled or suspended.

Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR USE IN WILDLIFE FOOD PLOTS IN WILDLIFE OPENINGS

Directions For Use:

Alligare IMOX Herbicide may be applied to crops listed in this label when they are planted solely for the use of wildlife food plots in wildlife openings. These species include alfalfa, beans (dry), chicory, clover grown for nonfood and nonfeed, clover grown for seed, edamame, lima bean (succulent), peas (dry), pea (English), snap bean, and soybean.

For applications to wildlife food plots, refer to weed control tables and restrictions for the applicable crop species on the IMOX Section 3 product label.

Areas that may be Grazed or Cut for Hay

Alligare IMOX Herbicide applied to listed aquatic and terrestrial non-crop sites may be grazed or cut for at a maximum use rate of 1 gallon per acre of Alligare Imox Herbicide or 5% v/v spray for spot treatments. There are no grazing or haying restrictions. Use Alligare IMOX Herbicide up to a rate of one gallon per year for selective weed control on forage intended for wildlife consumption in wildlife openings.

Terrestrial Use Directions

Alligare IMOX Herbicide can be applied as a foliar spot spray with the use of ground equipment in sufficient water to obtain uniform distribution and spray to targeted foliage.

Foliar Spot Application

Apply as a solution containing up to 5% v/v Alligare IMOX Herbicide spray.

RESTRICTIONS AND PRECAUTIONS

- DO NOT apply more than 1 pound of Imazamox acid equivalent (1 gallon) per acre per year.
- DO NOT exceed 2 applications of Alligare IMOX Herbicide per year.
- DO NOT harvest any plant species in wildlife food plots for seed, bedding, or human or livestock consumption.
- Minimum Retreatment Interval: 14 days

Alligare, LLC
1565 5th Avenue
Opelika, AL 36801

REV 121724